

The higher-order C_n dispersion coefficients for hydrogen.

J.Mitroy* and M.W.J.Bromley†

Faculty of Technology, Charles Darwin University, Darwin NT 0909, Australia

(Dated: February 2, 2008)

The complete set of 2nd, 3rd and 4th order van der Waals coefficients, C_n up to $n = 32$ for the H(1s)-H(1s) dimer are computed using pseudo-states to evaluate the appropriate sum rules. A study of the convergence pattern for $n \leq 16$ indicates that all the $C_{n \leq 16}$ coefficients are accurate to 13 significant digits. The relative size of the 4th-order $C_n^{(4)}$ to the 2nd-order $C_n^{(2)}$ coefficients is seen to increase as n increases and at $n = 32$ the 4th-order term is actually larger.

PACS numbers: 34.20.Cf, 31.25.Jf, 31.15.Pf, 32.10.Dk

I. INTRODUCTION

The long-range interaction between two spherically symmetric atoms can be written in the general form

$$V(R \rightarrow \infty) = -V_6(R) - V_8(R) - V_{10}(R) - V_{11}(R) - \dots, \quad (1)$$

where the dispersion potential, $V_n(R)$ of order n is written

$$V_n(R) = \frac{C_n}{R^n}. \quad (2)$$

The C_n parameters are the van der Waals dispersion coefficients. The even ($n = 6, 8, \dots$) dispersion coefficients are calculated using sum rules derived from 2nd-order perturbation theory and provide an attractive interaction. The odd ($n = 11, 13, \dots$) terms come from 3rd-order perturbation theory, and are repulsive [1, 2]. Contributions from 4th-order perturbation theory start at $n = 12$ [1, 3].

The dispersion interaction for the simplest system, namely the hydrogen dimer, is only known precisely to $n = 11$. The latest calculations by Yan and Dalgarno (YD) [2] reported almost exact values for the 2nd-order even dispersion parameters up to $n = 16$. They also gave almost exact results for the 3rd-order coefficients, up to $n = 15$. However, it is known that contributions from 4th-order perturbation theory start at $n = 12$ [1, 3, 4] so the dispersion potential computed from the YD C_n is incomplete for $n > 11$.

In this article, the complete set of dispersion parameters, from C_6 to C_{16} are computed by using a large pseudo-state expansion to evaluate the appropriate sum rules. The contributions from 4th-order perturbation theory to C_{12} , C_{14} and C_{16} are explicitly included.

II. CALCULATION OF THE DISPERSION PARAMETERS

All the dispersion coefficients computed in this paper were computed by first diagonalizing the semi-empirical Hamiltonian in a large basis of Laguerre type orbitals (LTOs) defined by

$$\chi_\alpha(r) = N_\alpha r^{\ell+1} \exp(-\lambda_\alpha r) L_{n_\alpha-\ell-1}^{(2\ell+2)}(2\lambda_\alpha r), \quad (3)$$

where the normalization constant is

$$N_\alpha = \sqrt{\frac{(2\lambda_\alpha)^{2\ell+3} (n_\alpha - \ell - 1)!}{(\ell + n_\alpha + 1)!}}. \quad (4)$$

The function $L_{n_\alpha-\ell-1}^{(2\ell+2)}(2\lambda_\alpha r)$ is an associated Laguerre polynomial that can be defined in terms of a confluent hypergeometric function [5] as

$$L_{n_\alpha-\ell-1}^{(2\ell+2)}(2\lambda_\alpha r) = \frac{(n_\alpha + \ell + 1)!}{(n_\alpha - \ell - 1)!(2\ell + 2)!} \times M(-(n_\alpha - \ell - 1), 2\ell + 2, 2\lambda_\alpha r) \quad (5)$$

All the matrix elements can be written in simple compact forms provided a common λ_α is adopted throughout the calculation. However, in the present work, the radial wave functions were placed on a numerical grid and all matrix elements computed by gaussian quadratures. This was done for reasons of convenience as the diagonalisation could be done with an existing program used in previous calculations of the dispersion parameters and the structures of positronic atoms [6, 7]. This program can achieve close to machine precision in almost all radial matrix computations.

Once the Hamiltonian diagonalisation is complete, sum rules involving radial matrix elements were used to determine the dispersion parameters. The specific sum-rules used are those derived by Ovsiannikov, Guilyarovski and Lopatko (OGL) [1]. Their expressions are a bit simpler than those developed by other authors [2, 3]. There were some omissions in their published equations [8], and a more thorough description of the sum-rules is presented here.

*Electronic address: jxm107@rsphysse.anu.edu.au

†Electronic address: mbromley@cdu.edu.au

TABLE I: The dispersion coefficients for the H-H dimer. All the results in the "Best Previous" column come from [2] except that for $C_{12}^{(4)}$ which is taken from [3]. All values are in atomic units.

Coefficient	10 LTOs	15 LTOs	20 LTOs	Best previous
C_6	6.499 026 705 3305	6.499 026 705 4057	6.499 026 705 4059	6.499 026 705 4058
C_8	124.399 083 58236	124.399 083 58362	124.399 083 58362	124.399 083 58362
C_{10}	3285.828 414 9425	3285.828 414 9674	3285.828 414 9674	3285.828 414 9674
$C_{12}^{(2)}$	$1.214\ 860\ 208\ 9619 \times 10^5$	$1.214\ 860\ 208\ 9686 \times 10^5$	$1.214\ 860\ 208\ 9686 \times 10^5$	$1.214\ 860\ 208\ 9686 \times 10^5$
$C_{14}^{(2)}$	$6.060\ 772\ 689\ 1671 \times 10^6$	$6.060\ 772\ 689\ 1917 \times 10^6$	$6.060\ 772\ 689\ 1917 \times 10^6$	$6.060\ 772\ 689\ 1921 \times 10^6$
$C_{16}^{(2)}$	$3.937\ 506\ 393\ 9865 \times 10^8$	$3.937\ 506\ 393\ 9985 \times 10^8$	$3.937\ 506\ 393\ 9985 \times 10^8$	$3.937\ 506\ 393\ 9992 \times 10^8$
C_{11}	-3474.898 037 8919	-3474.898 037 8822	-3474.898 037 8822	-3474.898 037 8822
C_{13}	$-3.269\ 869\ 240\ 4549 \times 10^5$	$-3.26\ 986\ 924\ 04407 \times 10^5$	$-3.26\ 986\ 924\ 04407 \times 10^5$	$-3.26\ 986\ 924\ 04407 \times 10^5$
C_{15}	$-2.839\ 558\ 063\ 3179 \times 10^7$	$-2.839\ 558\ 063\ 2998 \times 10^7$	$-2.839\ 558\ 063\ 2997 \times 10^7$	$-2.839\ 558\ 063\ 2998 \times 10^7$
$C_{12}^{(4)}$	1241.587 803 8317	1241.587 803 8462	1241.587 803 8462	1241.588
$C_{14}^{(4)}$	$3.009\ 633\ 558\ 9570 \times 10^5$	$3.009\ 633\ 559\ 0035 \times 10^5$	$3.096\ 633\ 559\ 0035 \times 10^5$	
$C_{16}^{(4)}$	$4.745\ 455\ 287\ 4168 \times 10^7$	$4.745\ 455\ 287\ 4083 \times 10^7$	$4.745\ 455\ 287\ 4079 \times 10^7$	
C_{12}	$1.227\ 276\ 087\ 0002 \times 10^5$	$1.227\ 276\ 087\ 0071 \times 10^5$	$1.227\ 276\ 087\ 0071 \times 10^5$	$1.227\ 27609 \times 10^5$ ^a
C_{14}	$6.361\ 736\ 045\ 0628 \times 10^6$	$6.361\ 736\ 045\ 0920 \times 10^6$	$6.361\ 736\ 045\ 0921 \times 10^6$	
C_{16}	$4.412\ 051\ 922\ 7282 \times 10^8$	$4.412\ 051\ 922\ 7393 \times 10^8$	$4.412\ 051\ 922\ 7393 \times 10^8$	

^aThis entry adds to BM $C_{12}^{(4)}$ to the YD $C_{12}^{(4)}$.

A. The 2nd-order terms

The 2nd-order dispersion coefficients for the H-H system have been determined to high accuracy [2, 9, 10] even for high n . The working expression adopted for computation is,

$$C_{2\lambda+6}^{(2)} = \sum_{\ell_1=1}^{\lambda+1} \frac{(2\lambda+4)!}{(2\ell_1+1)!(2\ell'_1+1)!} \times \sum_{i_1, i'_1} \frac{\langle 0, 0 | r^{\ell_1} | i_1, \ell_1 \rangle^2 \langle 0, 0 | r^{\ell'_1} | i'_1, \ell'_1 \rangle^2}{(E_{i_1} + E_{i'_1} - 2E_0)^2} \quad (6)$$

where $\ell'_1 = \lambda + 2 - \ell_1$. The state vector $|i_1, \ell_1\rangle^2$ represents the radial part of the state i_1 with orbital angular momentum ℓ_1 and energy E_{i_1} . The ground state energy is E_0 . The sum rule

$$T^{(\ell)} = \sum_i \langle 0, 0 | r^\ell | i, \ell \rangle^2 = \frac{(2\ell+2)!}{2^{(2\ell+1)}} \quad (7)$$

is a useful diagnostic check of the accuracy of the underlying discretization of the H-spectrum.

B. The 3rd-order terms; C_{11} and C_{13}

The dispersion coefficients, C_{11} and C_{13} , arise from 3rd-order perturbation theory [1, 2, 3, 4, 11]. Close to exact dispersion parameters for the H-H system have been published [2].

The general expression for the 3rd-order $C_{2\lambda+11}$ is [1]

$$C_{2\lambda+11} = - \sum_{\ell_1 k_1 \ell_2} \sum_{\ell'_1 k'_1 \ell'_2} \sum_{i_1 i'_1 i_2 i'_2} \frac{G(\lambda, \ell_1, \ell'_1, \ell_2, \ell'_2, k_1, k'_1)}{(E_{i_1 i'_1} - 2E_0)(E_{i_2 i'_2} - 2E_0)} \times \langle 0, 0 | r^{\ell_1} | i_1, \ell_1 \rangle \langle i_1, \ell_1 | r^{k_1} | i_2, \ell_2 \rangle \times \langle i_2, \ell_2 | r^{\ell'_2} | 0, 0 \rangle \langle 0, 0 | r^{\ell'_1} | i'_1, \ell'_1 \rangle \times \langle i'_1, \ell'_1 | r^{k'_1} | i'_2, \ell'_2 \rangle \langle i'_2, \ell'_2 | r^{\ell'_2} | 0, 0 \rangle \quad (8)$$

with the notation $E_{i_1 i'_1} = E_{i_1} + E_{i'_1}$ being used in the energy denominator. The parameter λ is defined

$$2\lambda + 8 = \ell_1 + k_1 + \ell_2 + \ell'_1 + k'_1 + \ell'_2 \quad (9)$$

and all of the angular momentum indices are greater than zero. Defining $J = (\ell_1 + k_1 + \ell_2)/2$ and $J' = (\ell'_1 + k'_1 + \ell'_2)/2$, the coefficient G is defined as

$$G(\lambda, \ell_1, \ell'_1, \ell_2, \ell'_2, k_1, k'_1) = (\lambda+4)! A(J, \ell_1, k_1, \ell_2) \times A(J', \ell'_1, k'_1, \ell'_2) B(\lambda, \ell_1, \ell'_1) B(\lambda, k_1, k'_1) B(\lambda, \ell_2, \ell'_2) \quad (10)$$

where

$$B(\lambda, \ell_1, \ell_2) = \frac{[2(\lambda+4-\ell_1-\ell_2)]!}{(\lambda+4-\ell_1-\ell_2)!}, \quad (11)$$

and

$$A(J, \ell_1, k_1, \ell_2) = \frac{J!}{(2J+1)!(J-\ell_1)!(J-k_1)!(J-\ell_2)!} \quad (12)$$

C. The 4th-order contributions to C_{12} and C_{14}

As far as we know, there have only been two explicit calculations of the 4th-order contribution to C_{12} . Bukta and Meath [3] gave estimates of $C_{12}^{(2)}$ and $C_{12}^{(4)}$ for the hydrogen dimer. Ovsinnikov *et al* [1] developed a general and compact expression for the evaluation of $C_n^{(4)}$, and in addition they reported values of $C_{12}^{(4)}$ for all possible combinations of hydrogen and the alkali atoms. Rectifying some omissions in their published equations [8], one writes

$$C_{2\lambda+12}^{(4)} = b_{2\lambda+12} - \sum_{\lambda_1=0}^{\lambda} C_{2\lambda_1+6}^{(2)} a_{2\lambda_2+6}, \quad (13)$$

where

$$\lambda = \lambda_1 + \lambda_2. \quad (14)$$

The factor $a_{2\lambda_2+6}$ is

$$a_{2\lambda_2+6} = \sum_{\ell_1=1}^{\lambda_2+1} \frac{(2\lambda_2+4)!}{(2\ell_1+1)!(2\ell'_1+1)!} \times \sum_{i_1, i'_1} \frac{\langle 0, 0 | r^{\ell_1} | i_1, \ell_1 \rangle^2 \langle 0, 0 | r^{\ell'_1} | i'_1, \ell'_1 \rangle^2}{(E_{i_1} + E_{i'_1} - 2E_0)^2} \quad (15)$$

where

$$\ell_1 + \ell'_1 = \lambda + 2. \quad (16)$$

The expression for $a_{2\lambda+6}$ is practically the same as eq. (6) for $C_{2\lambda+6}^{(2)}$; the only difference being an extra factor in the energy denominator (compare with eq. (10) of [1]).

The factor $b_{2\lambda+12}$ is more complicated and defined as

$$\begin{aligned} b_{2\lambda+12} = & \sum_{\ell_1, \ell_2, \ell_3} \sum_{\ell'_1, \ell'_2, \ell'_3} \sum_{k_1, k_2, k'_1, k'_2} \sum_K \sum_{i_1 i'_1 i_2 i'_2 i_3 i'_3} \\ & \times \left[\frac{\hat{L}_1! \hat{K}_1! \hat{K}_2! \hat{L}_3!}{2\ell_1! 2\ell'_1! 2k_1! 2k'_1! 2k_2! 2k'_2! 2\ell_3! 2\ell'_3!} \right]^{1/2} \\ & \times \langle \ell_1 0 k_1 0 | \ell_2 0 \rangle \langle \ell'_1 0 k'_1 0 | \ell'_2 0 \rangle \\ & \times \langle k_2 0 \ell_3 0 | \ell_2 0 \rangle \langle k'_2 0 \ell'_3 0 | \ell'_2 0 \rangle \\ & \times \langle L_1 0 K_1 0 | K 0 \rangle \langle K_2 0 L_3 0 | K 0 \rangle \\ & \times \begin{Bmatrix} \ell_1 & \ell'_1 & L_1 \\ k_1 & k'_1 & K_1 \\ \ell_2 & \ell'_2 & K \end{Bmatrix} \begin{Bmatrix} k_2 & k'_2 & K_2 \\ \ell_3 & \ell'_3 & L_3 \\ \ell_2 & \ell'_2 & K \end{Bmatrix} \\ & \times \frac{1}{(E_{i_1 i'_1} - 2E_0)(E_{i_2 i'_2} - 2E_0)(E_{i_3 i'_3} - 2E_0)} \\ & \times \langle 0, 0 | r^{\ell_1} | i_1, \ell_1 \rangle \langle i_1, \ell_1 | r^{k_1} | i_2, \ell_2 \rangle \\ & \times \langle i_2, \ell_2 | r^{k_2} | i_3, \ell_3 \rangle \langle i_3, \ell_3 | r^{\ell_3} | 0, 0 \rangle \\ & \times \langle 0, 0 | r^{\ell'_1} | i'_1, \ell'_1 \rangle \langle i'_1, \ell'_1 | r^{k'_1} | i'_2, \ell'_2 \rangle \\ & \times \langle i'_2, \ell'_2 | r^{k'_2} | i'_3, \ell'_3 \rangle \langle i'_3, \ell'_3 | r^{\ell'_3} | 0, 0 \rangle \end{aligned} \quad (17)$$

where $L_1 = \ell_1 + \ell'_1$, $L_3 = \ell_3 + \ell'_3$, $K_1 = k_1 + k'_1$ and $K_2 = k_2 + k'_2$. We use $\hat{L} = (2L + 1)$. The sums are constrained by the condition

$$L_1 + K_1 + K_2 + L_3 = 2\lambda + 8. \quad (18)$$

While $\ell_1, \ell'_1, \ell_3, \ell'_3$ must be greater than 0, it is possible for ℓ_2 and ℓ'_2 to be equal to 0. None of k_1, k'_1, k_2 or k'_2 can be zero. Since ℓ_2 and ℓ'_2 can both be equal to zero, the possibility of i_2, i'_2 both occupying the ground state must be explicitly excluded from the summation.

III. RESULTS OF THE CALCULATION

The results of the calculations for the complete set of dispersion coefficients up to C_{16} are given in Table I. The parameters are given for basis sets with 10, 15 and 20 basis functions per angular momentum respectively. The exponent in the LTO was chosen to be $\lambda = 1.0$ for all angular momenta. This choice resulted in much faster convergence of the dispersion parameters than that observed by Yan and Dalgarno in their calculations of the 3rd-order dispersion coefficients. Table I also gives results reported by YD and a single calculation of $C_{12}^{(4)}$ by Bukta and Meath (BM) [3].

A. The 2nd-order terms

The calculations of $C_n^{(2)}$ do not give new information and Yan and Dalgarno [2] have given values which are converged to better than 15 significant figures. The present calculations with the $N = 20$ basis are identical to 13 significant figures. The small differences in the last digit for some of coefficients arise from minor inaccuracies with the radial matrix elements. Hence we conclude that the present calculations are numerically reliable and that the pseudo-state representation of the H-spectrum is close to converged.

Besides the dispersion coefficients, the sum-rule, eq. (7) was evaluated and seen to be correct to 12 significant digits for all polarities relevant to the evaluation of C_6 to $C_{16}^{(2)}$.

B. The 3rd-order terms

Since the 3rd-order terms, $C_n^{(3)}$ have already been given by YD, these calculations merely serve as a test of our numerical procedures. Once again, calculations with the 20 LTO basis agree with the YD results to 14 significant figures. It is worth noting the present results required fewer terms than YD to achieve convergence. YD made the choice $\lambda_\alpha = 1/(\ell + 1)$ in eq. (3) and did not achieve convergence to the 14th digit place until the dimension of the LTO expansion was 50. The present basis with $\lambda_\alpha = 1.0$ achieves the same level of convergence with 20 LTOs.

TABLE II: The $n \geq 16$ dispersion coefficients for the H-H dimer. All values are in atomic units.

n	$C_n^{(2)}$	$C_n^{(4)}$	$C_n^{(2)} + C_n^{(4)}$	$C_n^{(3)}$
17				$2.726\ 099\ 889 \times 10^9$
18	$3.234\ 218\ 716 \times 10^{10}$	$7.009\ 061\ 179 \times 10^9$	$3.935\ 124\ 834 \times 10^{10}$	
19				$3.020\ 900\ 833 \times 10^{11}$
20	$3.278\ 573\ 440 \times 10^{12}$	$1.083\ 922\ 188 \times 10^{12}$	$4.362\ 495\ 628 \times 10^{12}$	
21				$3.900\ 227\ 980 \times 10^{13}$
22	$4.021\ 082\ 848 \times 10^{14}$	$1.832\ 218\ 347 \times 10^{14}$	$5.853\ 301\ 195 \times 10^{14}$	
23				$5.856\ 636\ 712 \times 10^{15}$
24	$5.868\ 996\ 335 \times 10^{16}$	$3.444\ 924\ 821 \times 10^{16}$	$9.313\ 921\ 156 \times 10^{16}$	
25				$1.017\ 059\ 252 \times 10^{18}$
26	$1.005\ 294\ 993 \times 10^{19}$	$7.249\ 737\ 286 \times 10^{18}$	$1.730\ 268\ 722 \times 10^{19}$	
27				$2.028\ 440\ 001 \times 10^{20}$
28	$1.996\ 944\ 941 \times 10^{21}$	$1.709\ 243\ 726 \times 10^{21}$	$3.706\ 188\ 667 \times 10^{21}$	
29				$4.613\ 037\ 362 \times 10^{22}$
30	$4.553\ 288\ 866 \times 10^{23}$	$4.507\ 006\ 859 \times 10^{23}$	$9.060\ 295\ 725 \times 10^{23}$	
31				$1.188\ 007\ 684 \times 10^{25}$
32	$1.181\ 107\ 088 \times 10^{26}$	$1.325\ 398\ 446 \times 10^{26}$	$2.506\ 505\ 534 \times 10^{26}$	

C. The 4th-order terms

The only previous explicit calculation of a 4th-order term was that made by Bukta and Meath (BM) [3], and the only parameter given was $C_{12}^{(4)}$. The OGL [1] estimate of $C_{12}^{(4)}$, 1.220×10^5 au was made using an approximation to the Greens function and so perfect agreement is not expected. However, the present calculation agrees with BM calculation of $C_{12}^{(4)}$ to all digits quoted, namely seven.

The number of terms in the sum, eq. (17) increases rapidly as n increases. There are 4 terms for $C_{12}^{(4)}$, there are 64 terms for $C_{14}^{(4)}$, and finally there are 460 terms for $C_{16}^{(4)}$.

The dominant contribution to $C_n^{(4)}$ comes from $b_{2\lambda+12}$ with 96% of $C_{12}^{(4)}$ coming from b_{12} . The tendency for $b_{2\lambda+12}$ to be the dominant term in $C_n^{(4)}$ becomes more accentuated as n increases and the b_{16} term gives an estimate to $C_{16}^{(4)}$ that is correct to 0.1%.

One feature of Table I concerns the relative size of $C_n^{(4)}$ to $C_n^{(2)}$. For $n = 12$, the $C_n^{(4)} : C_n^{(2)}$ ratio is 1.02%. However, as n gets larger, the ratio also gets larger. For $n = 14$ the ratio is 4.97%, while for $n = 16$ the ratio is 12.1%.

D. The dispersion coefficients for $n \geq 17$

Higher order contributions than 4th-order begin at $n = 17$. There is a 5th-order contribution to C_{17} and a 6th-order contribution to C_{18} [1]. Estimates of $C_n^{(2)}$ for $n \geq 17$ have been made by a variety of authors [1, 10, 12, 13, 14]. However, the only estimate of the 3rd

and 4th-order terms with $n \geq 17$ are those of OGL [1]. By explicit calculation they obtained $C_{17}^{(3)} = -2.739 \times 10^9$ au which agrees with the present more extensive calculation to within 1%. Making an approximation to the greens function they estimated $C_{18}^{(4)} = 3.3 \times 10^9$ au, which is about half the size of the present value.

The dispersion parameters up to C_{32} from the present calculation are tabulated in Table II. The reason for taking the calculations so far rests in the relative size of the $C_n^{(4)}$ and $C_n^{(2)}$ terms. It was noticed that the $C_n^{(4)} : C_n^{(2)}$ ratio got larger as n increased. So the calculations were extended to C_{32} in order to demonstrate explicitly that the $C_n^{(4)} : C_n^{(2)}$ ratio can actually become larger than 1.0.

The precision of the entries in Table II is not as high as those in Table I. The calculations of $C_n^{(4)}$ did become more time consuming as n increased. There were 922,064 different $(\ell_1, k_1, \ell_2, \dots)$ combinations by the time $n = 32$ was reached. Also the number of radial integrals in eq. (17) increases as N^6 where N is the numbers of LTOs for any given ℓ . So the $N = 20$ calculation is 64 times more intensive than the $N = 10$ calculation.

The $C_n^{(2)}$ and $C_n^{(3)}$ entries in Table II were taken from the calculation with 15 LTOs. The $C_n^{(4)}$ entries were taken from a $N = 15$ calculation up to $n = 20$, thereafter the $N = 10$ basis was used. The values of $C_n^{(2)}$ agree with those of Thakkar [10] for all ten digits given in Table II. Comparisons between $N = 10$ and $N = 15$ calculations for $C_n^{(3)}$ suggest that the convergence is slower as n increases and that $C_{31}^{(3)}$ is reliable to about 6 digits. A similar level of accuracy can be expected for $C_n^{(4)}$ and a comparison between the $N = 10$ and $N = 15$ values for $C_{20}^{(4)}$ gives agreement for the first 9 digits.

IV. CONCLUSIONS

The higher n dispersion parameters from C_{11} , through to C_{16} have been computed to an accuracy of 13 significant figures for the H-H dimer. Since the 4th-order contributions were included for C_{12} , C_{14} and C_{16} the adiabatic dispersion interaction can now be regarded as complete up to terms of order R^{-16} .

The time taken to evaluate the dispersion coefficients was not excessive. For example, a calculation using 20 LTOs took about 17 minutes to determine all terms up to C_{16} on a 850 MHz CPU. Hence the pseudo-state method adopted here, and in other similar works (e.g. [2, 3]), could be used to make explicit calculations of the 5th-order correction to C_{17} and even the 6th-order correc-

tion to C_{18} [1]. Therefore, it is certainly possible with existing technology to determine the complete dispersion interaction for the H-H interaction for all terms up to and including C_{22} .

V. ACKNOWLEDGMENTS

The authors would like to thank Mr J C Nou and Mr C Hoffman of CDU for workstation support and Professor V D Ovsiannikov of Voronezh University for helpful communications about the exact form of the 4th-order matrix element and for pointing out some faults with this manuscript.

-
- [1] V. D. Ovsiannikov, A. V. Guilyarovski, and O. Y. Lopatko, *Mol. Phys.* **61**, 111 (1988).
 - [2] Z. C. Yan and A. Dalgarno, *Mol. Phys.* **96**, 863 (1999).
 - [3] J. F. Bukta and W. J. Meath, *Mol. Phys.* **27**, 1235 (1974).
 - [4] A. Dalgarno, *Adv. Chem. Phys.* **12**, 143 (1967).
 - [5] M. Abramowitz and I. E. Stegun, eds., *Handbook of Mathematical Functions* (US GPO, Washington DC, 1972), Natl. Bur. Stand. Appl. Math. Ser. 55.
 - [6] M. W. J. Bromley and J. Mitroy, *Phys. Rev. A* **65**, 012505 (2002).
 - [7] J. Mitroy and M. W. J. Bromley, *Phys. Rev. A* **68**, 052714 (2003).
 - [8] V. D. Ovsiannikov, priv. comm. (2004).
 - [9] W. J. Deal, *Int. J. Quantum Chem.* **6**, 593 (1972).
 - [10] A. J. Thakkar, *J. Chem. Phys.* **89**, 2092 (1988).
 - [11] G. P. Arrighini, F. Biondi, and C. Guidotti, *Mol. Phys.* **26**, 1137 (1973).
 - [12] R. J. Bell, *Proc. Phys. Soc. London* **86**, 239 (1965).
 - [13] R. J. Bell, *Proc. Phys. Soc. London* **87**, 594 (1966).
 - [14] A. Koide, W. J. Meath, and A. R. Allnatt, *Chem. Phys.* **58**, 105 (1981).